

This product cleans by removing dirt, grime, blood, urine, fecal matter and other common soils found in animal housing facilities, grooming facilities, kennels, pet stores, veterinary clinics, laboratories or other small animal facilities.

This product can be used on washable hard, non-porous surfaces such as: glass surfaces, laminated surfaces, metal, stainless steel, glazed porcelain, glazed ceramic, sealed granite, sealed marble, plastic such as polystyrene or polypropylene, sealed limestone, sealed slate, sealed stone, sealed terra cotta, sealed terrazzo, chrome, vinyl, Plexiglass®.

**DISINFECTION PERFORMANCE IN INSTITUTIONS AND NON-MEDICAL FACILITIES:** This product is effective against: *Pseudomonas aeruginosa*, *Salmonella enterica*, *Staphylococcus aureus*

**VIRUCIDAL\* PERFORMANCE:** This product is effective against: Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), Herpes Simplex virus Type 1, Human Coronavirus, HIV-1, Influenza A2/Japan virus, Influenza A H1N1 Virus, Norovirus

**ANIMAL VIRUCIDAL\* PERFORMANCE:** This product is effective against: Avian Reovirus, Bovine Viral Diarrhea virus, \*\*Canine Adenovirus (1.25 oz/1 gal. of water), Canine Coronavirus, Canine Distemper virus, \*\*Canine Parainfluenza Virus (1.25 oz/1 gal. of water), \*\*\*Canine Parvovirus (CPV) (4 oz/1 gal. of water), Equine Arteritis Virus, \*\*Feline Calicivirus (1.25 oz/1 gal. of water), \*\*Feline Infectious Peritonitis virus (1.25 oz/1 gal. of water), \*\*Feline Leukemia virus (1.25 oz/1 gal. of water), \*\*Feline Panleukopenia virus (1.25 oz/1 gal. of water), \*\*Feline Picornavirus (1.25 oz/1 gal. of water), \*\*Feline Rhinotracheitis virus (1.25 oz/1 gal. of water), Infectious Avian Laryngotracheitis virus, Infectious Bovine Rhinotracheitis virus (IBR), Infectious Bronchitis Virus, \*\*Infectious Canine Hepatitis Virus (1.25 oz/1 gal. of water), \*\*\*Minute virus of Mice Parvovirus (4 oz/1 gal. of water), Murine Norovirus, \*\*Murine Parainfluenza virus type 1 (1.25 oz/1 gal. of water), Newcastle disease virus, \*\*\*Porcine Parvovirus (4 oz/1 gal. of water), Porcine Respiratory & Reproductive Virus (PRRSV), Porcine Rotavirus, Pseudorabies virus, \*\*Rabies virus (1.25 oz/1 gal. of water), Transmissible Gastroenteritis virus (TGE), Vaccinia virus

**FUNGICIDAL PERFORMANCE:** This product is effective against: *Trichophyton mentagrophytes*

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Before use in federally inspected meat and poultry food processing plants and dairies, food products and packaging materials must be removed from room or carefully protected. Prior to use of this product, cover or remove all food and packaging material before disinfection and remove all animals and feeds from areas to be treated, animal transportation vehicles, and enclosures. Remove all litter, droppings and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other surfaces of facilities and fixtures occupied or traversed by poultry or other animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean surfaces with soap or detergent and rinse with water. **Do not use in milking stalls, milking parlors or milk houses.**

**DISINFECTION/VIRUCIDAL\*/FUNGICIDAL/PERFORMANCE IN INSTITUTIONS AND NON-MEDICAL FACILITIES:** Add \*\*\*1.25 oz of this product to 1 gallon of water. For heavily soiled areas, a preliminary cleaning is required. Apply use solution to hard non-porous surfaces with a brush, cloth, mop, sponge, auto scrubber, mechanical spray device or by immersion so as to thoroughly wet surfaces. For sprayer applications, spray 6-8 inches from surface. Do not breathe spray.

Immerse all halters and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure in the use solution. After application, ventilate buildings, coops and other closed spaces. Do not house animals, or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, troughs, automatic feeders, fountains and waterers and other treated equipment which can contact food or water with soap or detergent, and rinse with potable water before reuse.

Rub with brush, cloth, mop, or sponge. Treated surfaces must remain wet for 10 minutes. Allow to air dry. Rinsing of floors is not necessary unless they are to be waxed or polished. Change cloth, sponge or towels frequently to avoid redeposition of soil. Prepare a fresh solution at least daily or when use solution becomes visibly dirty.

**FOR DISINFECTING HARD, NON-POROUS SURFACES AGAINST INFLUENZA A H1N1 VIRUS:** For use in hospitals, animal premises, and other non-animal use sites, follow the **DISINFECTION/ VIRUCIDAL\*/FUNGICIDAL/PERFORMANCE IN INSTITUTIONS AND NON-MEDICAL FACILITIES** directions.

**HOSPITAL/HEALTH CARE/INSTITUTIONAL**

This product is not for use on Medical device surfaces.

**\*KILLS HIV, HCV and HBV ON PRE-CLEANED ENVIRONMENTAL SURFACES/ OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of hard non-porous surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type 1 (HIV-1), Hepatitis B virus (HBV) and Hepatitis C virus (HCV).

**VIMOB A 128**

**CLEANER, DISINFECTANT, FUNGICIDE, VIRUCIDE\***  
With Organic Soil Tolerance For Institutional, Industrial, Veterinary, Federally Inspected Meat and Poultry Plants.

**ACTIVE INGREDIENTS:**

Alkyl (C14 60%, C16 30%, C12 5%, C18 5%)	4.5%
dimethyl benzyl ammonium chloride	
Alkyl (C12 68%, C14 32%)	4.5%
dimethyl ethylbenzyl ammonium chloride	
<b>INERT INGREDIENTS:</b>	<b>91.0%</b>
<b>TOTAL:</b>	<b>100.0%</b>

**KEEP OUT OF REACH OF CHILDREN  
DANGER PELIGRO**

See side panel for additional precautionary statements.  
**FIRST AID**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

EPA Reg. No. 10324-58-46269

EPA Est. No. 46269-DE-01

**NET CONTENTS:**

☐ 15 gallons ☐ 30 gallons ☐ 55 gallons

LOT #: **VM128092211**



**B14**

**QUIP LABORATORIES, INC.**  
1500 EASTLAWN AVENUE • WILMINGTON, DE 19802  
TEL : 800-424-2436 FAX: 302-761-2611

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HCV and HBV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.**

**Personal Protection:** Clean up must always be done wearing protective latex gloves, gowns, masks and eye protection. **Cleaning Procedure:** Blood and other body fluids must be thoroughly cleaned from hard non-porous surfaces and objects before application of this product. **Disposal of Infectious Materials:** Blood and other body fluids, cleaning materials and clothing must be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal. **Contact Time:** For HIV only, leave hard nonporous surfaces wet for 2 minutes. For all other viruses and bacteria, leave hard nonporous surfaces wet for 10 minutes. The 2-minute contact time will not control other common types of viruses and bacteria listed on this label.

**EFFECTIVENESS AGAINST CANINE PARVOVIRUS (CPV), MINUTE VIRUS OF MICE PARVOVIRUS AND PORCINE PARVOVIRUS:** Saturate surfaces with the disinfecting solution of 4 ounces per 1 gallon of water (or equivalent use dilution) for a period of 10 minutes. Immerse all halters, ropes and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure. After application, ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. All treated equipment that will contact feed or drinking water must be thoroughly scrubbed with soap or detergent then rinsed with potable water before reuse. Change cloth, sponge or towels frequently to avoid redeposition of soil. Prepare a fresh solution at least daily or when use solution becomes visibly dirty.

**HOSE DISPENSER:** (Note: this is to be used with the above claim only): The hose end sprayer is calibrated to deliver 4 ounces per gallon of water, and when hose is attached to this bottle will treat 12,000 square feet. Attach sprayer to hose. Secure tightly. Check that the sprayer is in the off position. Turn on water. Turn sprayer to on position. Spray evenly over surface. When finished turn sprayer to off position and then turn water off. Separate mixing of the concentrate or other application equipment is not required. First ensure the hose faucet is turned off.

**FOR USE AS A GENERAL CLEANER:** Apply use solution of 1-2 ounces of this product per 1 gallon of water to surfaces. Allow to air dry.

**GLASS CLEANING DIRECTIONS:** Use a 1-2 ounce of this product per 1 gallon of water use solution to clean windows, mirrors, and glass surfaces. Use a coarse spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray. Rub with sponge or cloth. Change cloth, sponge or towels frequently to avoid redeposition of soil.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**DANGER.** Corrosive. Causes irreversible eye damage and skin burns. Harmful if swallowed or absorbed through the skin. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves and protective clothing when handling. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish and aquatic invertebrates. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product into sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

**PHYSICAL OR CHEMICAL HAZARDS**

Do not use or store near heat or open flame. Do not mix oxidizers, anionic soaps and detergents.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage and disposal. **PESTICIDE STORAGE:** Store only in original container. Keep this product under locked storage sufficient to make it inaccessible to children or persons unfamiliar with its proper use.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

**CONTAINER HANDLING:** Non-Refillable Container. Do not reuse or refill this container. Triple rinse container promptly after emptying. Triple rinse as follows: Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip back and forth several times. Turn the container over onto its other end and tip back and forth several times. Follow Pesticide Disposal instructions for rinsate disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

**APPROVED**

*By Cherita Holt at 8:02 am, Mar 20, 2015*