

HEXAGIENE™

One-Step¹ Disinfectant²/Cleaner/Tuberculocide/Deodorizer (Meets OSHA Blood borne Pathogen Standard for *HBV, *HCV & *HIV)

ACTIVE INGREDIENTS

Active ingredient:	
Hydrogen peroxide.....	1.24%
Other ingredients:.....	98.76%
Total.....	100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

See Back Panel for Precautionary Statements and First Aid

Kills:

Bacteria: *Pseudomonas aeruginosa* (ATCC 15442), *Salmonella enterica* (ATCC 10708), *Staphylococcus aureus* (ATCC 6538), *Acinetobacter baumannii* (ATCC BAA-1709), *Enterobacter aerogenes* (ATCC 13048), (New Delhi metallo-beta-lactamase 1 (NDM-1) producing) *Enterobacter cloacae* (CDC 1000654), *Vancomycin Resistant Enterococcus faecalis* (VRE) (ATCC 51575), *Enterococcus faecium* (ATCC 6569), *Escherichia coli* (ATCC 11229), *Escherichia coli* O157:H7 (ATCC 35155), *Klebsiella pneumoniae* (ATCC 4352), *Carbapenem Resistant Klebsiella pneumoniae* (CRE) (K. pneumoniae) (ATCC BAA-1705), (New Delhi metallo-beta-lactamase 1 (NDM-1) producing) *Klebsiella pneumoniae* (CDC 1000527), *Vancomycin Resistant Staphylococcus aureus* (VRSA) (NARSA VR51), *Methicillin Resistant Staphylococcus aureus* (MRSA) (ATCC 33592), *Mycobacterium* (TB) *Mycobacterium bovis* BCG (TB) (ATCC 35743)

Viruses*: *Adenovirus Type 5 (ATCC VR-5) (Strain Adenoid 75), *Hepatitis B (HBV), *Hepatitis C (HCV) (ATCC VR-1422) (BVDV), *Herpes simplex virus type 1 (ATCC VR-733) (Strain F(1)), *HIV-1 (Strain HTLC-IIIb), *Human Rotavirus (ATCC VR-2018) (Strain WA), *Influenza A Virus (ATCC VR-544) (Strain A/8/68), *Norovirus (*Norwalk Virus) (ATCC VR-782), *SARS-Related Coronavirus 2 (cause of COVID-19), *Rhinovirus type 37 (ATCC VR-1147) (Strain 151-1)

Animal Viruses*: *Canine Parvovirus (ATCC VR-2017) (Strain Cornell-780916-80) in 5 minutes, *Feline Calicivirus (ATCC VR-782), *Minute virus of mice [ATCC VR-1346] (Strain Prototype (p))

Fungi: *Candida auris* (BAR-BANK#0381 from CDC)

Reduce cross contamination between treated surfaces¹. Use on Surfaces¹ in Food Storage Areas, Ambulatory Care Centers, Ambulatory Surgical Centers, Anesthesia Rooms, Birthing Center, Cat Lab, Central Service Areas, Central Supply Rooms, Critical Care Units, Dental Offices, Dialysis Clinics, Doctor's Offices, Donation Centers, Examination Rooms Or Areas, Emergency Rooms, Eye Surgical Centers, Home Health Care, Hospices, Hospital Beds, Hospital Environmental Services (EVS), Hospitals, ICU Areas, Isolation Areas, Laboratories, Long Term Care Facilities, Clinics, Medical Offices, Neonatal Intensive Care Units (NICU), New Born Nursery, Nursing Homes, Nursing's Stations, Skilled Nursing Facilities, Operating Rooms, Ophthalmic Offices, Orthopedics, Out-patient Surgical Centers (OPSC), Patient Care Areas, Patient Rooms, Pediatric Intensive Care Units (PICU), Pharmacies, Physician's Offices, Psychiatric Facilities, Public Care Areas, Radiology X-ray Rooms, X-ray Equipment, Recovery Rooms, Rehabilitation Centers, Respiratory Centers, Respiratory Therapy Rooms, Sick Rooms, Surgery Rooms, Surgical Centers, Surgery and Surgical Intensive Care Unit (SICU), Veterinary Clinics, and Waiting Rooms.

Use Hexagiene In: Institutional Facilities and Universities.

Hexagiene kills canine parvovirus and feline calicivirus on environmental surfaces¹ and kills *HIV-1 on environmental surfaces¹ in just 1 minute².

Use this product on washable, hard, nonporous surfaces¹ of: Autoclaves, Cabinet Handles, Computer Peripherals, Diagnostic Equipment, Equipment Surfaces, Equipment Tables, Exam Tables, High Touch Surfaces, Infant Incubators, Instrument Stands, Lab Benches, Medical Scales, Microscope, Rescue Tools, Stethoscopes, Other hard nonporous surfaces made of: Acrylic, Formica[®], Glass, Glazed Ceramic, Glazed Enameled Surfaces, Glazed Porcelain, Metal, Plastic such as Polystyrene or Polypropylene, Stainless Steel, Upholstery-vinyl and Plastic, Power Cords, PVC Tubing, Remote Controls, Thermometers, Transport Monitors, Ultra-sound Transducers And Equipment, Walls, Animal Equipment, Animal Crates, Automatic Feeders, Buckets, Brushes, Cages, Counters, Desks, Dog Runs, Examination Tables, External Surfaces of Veterinary Equipment Watering Appliances, Pens, Stalls, Transportation Cages.

Formica[®] is a registered trademark of the Diller Corporation

EPA REG. NO. 6836-442-46269 EPA EST. NO. 46269-DE-01

Net Contents: 1 Quart

Directions for use:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Unscrew cap and transfer your trigger to this bottle. This product is not to be used as a terminal sterilant / high level disinfectant on any surface or treatment that is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

To Use as a One-Step¹ Disinfectant²:

1. Pre-clean visibly soiled areas. 2. Spray 6 – 8 inches from surface¹, making sure to wet thoroughly. 3. Allow surface¹ to remain wet for 1 minute². For SARS-CoV-2, treated surfaces¹ must remain visibly wet for 30 seconds. 4. Wipe surface dry. 5. Allow to air dry.

To Disinfect Non-Critical, Pre-cleaned Instruments:

1. Instruments must be thoroughly pre-cleaned to remove excess organic debris, rinsed, and rough dried. (Clean and rinse lumens of hollow instruments before spraying with this product).

2. Spray all surfaces¹ of instruments with this product until thoroughly wet. 3. Allow surface¹ to remain wet for 1 minute². 4. Wipe with a clean, damp cloth or paper towel and allow to air dry. **Note:** Non-critical medical items are those that only come into contact with intact skin.

To Disinfect Surfaces¹ in Pharmaceutical and Cosmetic Processing Facilities:

1. Pre-clean visibly soiled areas. 2. Spray 6-8 inches from surface¹ until surface is thoroughly wet. 3. Allow surfaces to remain wet for 1 minute². 4. Rinse with sterile water. 5. Wipe with a dry, clean cloth and let air dry.

***KILLS HIV ON PRE-CLEANED ENVIRONMENTAL SURFACES¹ / OBJECTS PREVIOUSLY SOILED WITH BLOOD / BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces / objects with blood or body fluids, and in which the surfaces / objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 [HIV-1] [associated with AIDS], Hepatitis C Virus (HCV) or Hepatitis B Virus (HBV).

Special Instructions for Cleaning and Decontamination against HIV-1 on Surfaces / Objects Soiled With Blood / Body Fluids. Personal protection: Clean-up must always be done wearing protective gloves, gowns, masks and eye protection. **Cleaning procedure:** Blood and other body fluids containing HIV must be thoroughly cleaned from surfaces and objects before application of Hexagiene. Hexagiene can be used for this purpose. **Contact time:** Leave surface wet for 1 minute for HIV-1. Use appropriate contact time as listed on label for all other viruses, and bacteria claimed. **Disposal of infectious material:** Blood, body fluids, cleaning materials and clothing must be disposed of according to local regulations for infectious waste disposal.

Animal Housing Facilities including Veterinary Clinics, Equine Farms, Livestock, Swine and Poultry Houses: For cleaning and disinfecting the following surfaces¹: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries, etc. 1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc. 2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. 3. Empty or cover all troughs, racks, and other feeding and watering appliances. 4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water. 5. Saturate all surfaces by spraying with product until thoroughly wet. To disinfect, all surfaces must remain wet for 1 minute.² 6. Immerse or spray all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure. 7. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. 8. Thoroughly scrub treated feed racks, troughs, mangers, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

For use as a One-Step Cleaner and Disinfectant by Electrostatic Spraying: Ensure bystanders and pets are not present in the room during application. Clean visibly dirty surfaces prior to spraying. Wear a half face respirator with a chemical specific cartridge and a N95 filter, and eye protection (goggles, face shield or safety glasses) when spraying. Select a spray nozzle that produces spray droplet particles with a volume median diameter (VMD) ≥80 µm

Special Instructions for Cleaning Prior to Disinfection against Candida auris

Personal Protection: Wear appropriate barrier protection such as gloves, gowns, masks, or eye covering. **Cleaning Procedure:** Fecal matter/waste must be thoroughly cleaned from surfaces/objects before disinfection by application with a clean cloth, mop, and/or sponge saturated with the product. Pre-cleaning is to include vigorous wiping and/or scrubbing and all visible soil is removed. Surfaces in patient rooms are to be cleaned in an appropriate manner, such as from right to left or left to right, on horizontal surfaces, and top to bottom, on vertical surfaces, to minimize spreading the organism. Restrooms are to be cleaned last. Do not reuse soiled cloths. **Infectious Waste Disposal:** Materials used in the cleaning process that may contain feces/wastes are to be disposed of immediately in accordance with local regulations for infectious materials disposal.

Storage and Disposal:

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Open dumping is prohibited. Store in original container in areas inaccessible to children. **Container Disposal:** Nonrefillable container. Do not reuse or refill this container. **Plastic Containers:** Triple rinse (or equivalent). Offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID: IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a Poison Control Center or doctor for treatment advice. **IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a Poison Control Center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor or going for treatment.

¹ On hard, non-porous, non-food surfaces; not for indirect or direct food contact. ² 5-minute contact time is required for Canine Parvovirus efficacy claim.

³ When directions for disinfection are followed

MANUFACTURED FOR:



1500 EASTLAWN AVE. WILMINGTON DE. 19802
(800) 424-2436 | WWW.QUPLABS.COM