

Effective in hard water up to 400 ppm hardness calculated as Ca CO₃ in the presence of 5% soil. This product is for use on hard, non-porous surfaces in: Veterinary clinics, animal life science laboratories, animal laboratories, animal research centers, animal quarantine areas, animal holding areas, dog/cat animal kennels, pet animal quarters, pet shops, operating rooms, washing areas, waiting rooms, examination rooms and other animal care facilities.

This product can be used on washable hard, non-porous surfaces such as: Kennel runs, cages, kennel/cage floors, conductive flooring, veterinary x-ray tables, glazed porcelain, glazed tile, glazed ceramic, plastic, chrome and vinyl, enameled surfaces, painted surfaces, stainless steel, brass, copper, floors, walls, ceilings, fixtures, and countertops.

This product is effective against Canine Parvovirus**.

DISINFECTION PERFORMANCE: This product kills the following bacteria in 10 minutes at 1 oz. per gal. of water and 5% soil on hard, non-porous surfaces: Pseudomonas aeruginosa, Salmonella enterica, Staphylococcus aureus

VIROCIDAL* PERFORMANCE: This product kills the following viruses in 10 minutes at 1 oz. per gal. of water and 5% soil on hard, non-porous surfaces: Avian Influenza A (H5N1) Virus, Avian Influenza A (H3N2) Virus, Cytomegalovirus, Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), Herpes Simplex Virus Type 1, Herpes Simplex Virus Type 2, Human Coronavirus, Human Immunodeficiency Virus Type 1 (HIV 1), Influenza A Virus, Respiratory Syncytial Virus

ANIMAL PREMISE VIROCIDAL* PERFORMANCE: This product kills the following viruses in 10 minutes at 1 oz. per gal. of water and 5% soil on hard, non-porous surfaces: Avian Influenza A Virus (H5N1), Avian Influenza A Virus (H3N2), Canine Distemper Virus, Feline Picornavirus, Infectious Bovine Rhinotracheitis Virus, Pseudorabies Virus, Transmissible Gastroenteritis Virus

This product kills the following viruse in 10 minutes at 4 oz. per gal. of water and 5% soil on hard, non-porous surfaces: Canine Parvovirus**

** Indicates that a dilution of 4 oz. per gal. of water is required for this claim.

FUNGICIDAL PERFORMANCE: This product kills the following fungi in 10 minutes at 1 oz. per gal. of water and 5% soil on hard, non-porous surfaces: Candida albicans, Trichophyton mentagrophytes

MILDEWSTATIC PERFORMANCE: This product controls the following mold in 10 minutes at 1 oz. per gal. of water on hard, non-porous surfaces: Aspergillus niger

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR USE AS A ONE-STEP, GENERAL DISINFECTANT, FUNGICIDE, VIROCIDE*, CLEANER:

1. Pre-clean heavily soiled areas.
2. Apply use solution of 1 oz. of this product per gal. of water to disinfect hard, non-porous surfaces with a sponge, brush, cloth, mop, by immersion, auto scrubber, mechanical spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray.
3. Treated surfaces must remain wet for 10 minutes.
4. Allow to air dry.
5. Prepare a fresh solution daily or when visibly dirty.

***KILLS HIV, HBV AND HCV ON PRE-CLEANED HARD, NON-POROUS SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings or other settings in which there is an expected likelihood of soiling of hard, non-porous surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1), Hepatitis B Virus (HBV) and Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV and HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS. Personal Protection: Wear protective latex gloves, gowns, masks and eye protection. **Cleaning Procedure:** Blood and other body fluids containing HIV-1, HBV & HCV must be thoroughly cleaned from hard, non-porous surfaces and objects before application of this product. **Disposal of Infectious Materials:** Blood and other body fluids, cleaning materials and clothing must be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal. **Contact Time:** Allow hard, non-porous surfaces to remain wet for 10 minutes for all organisms.

TO CONTROL MOLD/MILDEW: Pre-clean heavily soiled areas. Prepare use solution by adding 1 oz. of this product per gal. of water. Apply use solution to hard, non-porous surfaces for 10 minutes which will effectively inhibit the growth of mold and mildew and their odors. Repeat treatment every seven days, or more often if new growth appears.

SANI-PLEX 128M

One-Step Disinfectant Cleaner

Disinfectant, Cleaner, Fungicide, Mildewstat, Virucide*. For use in Commercial Sites, Institutional Sites, Industrial Sites, and Veterinary Clinics.

ACTIVE INGREDIENTS:	
Didecyl Dimethyl Ammonium Chloride.....	5.07%
Alkyl (50% C14, 40% C12, 10% C16)	
Dimethyl Benzyl Ammonium Chloride.....	3.38%
OTHER INGREDIENTS:	91.55%
TOTAL:	100.00%

KEEP OUT OF REACH OF CHILDREN

DANGER PELIGRO

See side panel for additional precautionary statements.

FIRST AID

In case of emergency, call a poison control center or doctor for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

IF SWALLOWED: Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. Call a poison control center or doctor immediately for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

SI USTED NO ENTIENDE LA ETIQUETA, BUSQUE A ALGUIEN PARA QUE SE LA EXPLIQUE A USTED EN DETALLE. IF YOU DO NOT UNDERSTAND THE LABEL, FIND SOMEONE TO EXPLAIN IT TO YOU IN DETAIL.

EPA Reg. No. 10324-155-46269

EPA Est. No. 46269-DE-01

NET CONTENTS: 1 Gallon

B16



Quip Laboratories, Inc.

1500 EASTLAWN AVENUE, WILMINGTON, DE 19802

TEL: 302-761-2600 FAX: 302-761-2611

ANIMAL PREMISES: Prior to use of this product, remove all animals and feeds from areas to be treated, animal transportation vehicles (trucks, cars), and enclosures (coops, crates, kennels, stables). Remove all litter, droppings and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other surfaces of facilities and fixtures occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean surfaces with soap or detergent and rinse with water.

FOR USE AS AN ANIMAL PREMISE DISINFECTANT/VIROCIDE*: Prepare a use solution of 1 oz. of this product per gal. of water (1:128) (660 ppm active) for effectiveness against designated animal viruses on the label.

FOR USE AS AN ANIMAL PREMISE DISINFECTANT/VIROCIDE:** Prepare a use solution of 4 oz. of this product per gal. of water (1:32) (2600 ppm active) for effectiveness against designated animal viruse on the label.

Apply use solution to disinfect hard, non-porous surfaces with a sponge, brush, cloth, by immersion, mechanical spray device. For spray applications, spray 6-8 inches from surface. Do not breathe spray. Immerse all halters and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure in the use solution. Treated surfaces must remain wet for 10 minutes. Ventilate buildings, coops and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, troughs, automatic feeders, fountains and waterers and other treated equipment which can contact food or water with soap or detergent, and rinse with potable water before reuse.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store only in original container. Keep this product under locked storage sufficient to make it inaccessible to children or persons unfamiliar with its proper use.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Non-Refillable Container. Do not reuse or refill this container. Triple rinse container promptly after emptying. Triple rinse as follows: Fill the container ¼ full with water and recap. Shake for 10 seconds. Drain for 10 seconds after the flow begins to drip. Follow Pesticide Disposal instructions for rinsate disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER. Corrosive. Causes irreversible eye damage and skin burns. Harmful if swallowed or absorbed through the skin. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves and protective clothing when handling. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, oysters and shrimp.

