

VIMOB^A128

Efficacy as a Disinfectant

For animal water systems | add CDG Solution 3000 to water at a minimum dose of .5 ppm and a maximum dose of 5.0 ppm (5.0 mg/L) chlorine dioxide (using a dilution ratio of 1:600). For hard surface disinfection | please refer to label for dosing.

Type	Organism	Contact Time	Dilution
Disinfection	Acinetobacter baumannii Enterococcus faecalis Escherichia coli Klebsiella pneumoniae Pseudomonas aeruginosa Salmonella enterica Staphylococcus aureus	10 minutes	3 fl. oz. per gal. of 400 ppm hard water (2100 ppm active)
Virucidal	Avian Influenza A Herpes Simplex Virus Type 1 Herpes Simplex Virus Type 2 Human Immunodeficiency Virus Type 1 Human Coronavirus	10 minutes	3 fl. oz. per gal. of 400 ppm hard water (2100 ppm active)
Animal Premise Virucidal	Avian Influenza A Newcastle Disease Virus Transmissible Gastroenteritis Virus	10 minutes	3 fl. oz. per gal. of 400 ppm hard water (2100 ppm active)
Food Contact Surface Sanitizing	Escherichia coli Staphylococcus aureus	1 minute	1.25 fl. oz. per 3.5 gal. of 400 ppm hard water (250 ppm active)
Non-Food Contact Surface Sanitizing	Klebsiella pneumoniae Staphylococcus aureus	3 minutes	3 fl. oz. per gal. of 400 ppm hard water (2100 ppm active)
Fungicidal	Trichophyton interdigitale	10 minutes	3 fl. oz. per gal. of 400 ppm hard water (2100 ppm active)
Mildewstatic	Aspergillus niger	7 days	3 fl. oz. per gal. of 400 ppm hard water (2100 ppm active)